


Sample Name:
 ω -Hydroxy Terminated Polystyrene

Sample #: P18845- SOH

Structure:



Composition:

Mn x 10 ³	PDI
8.5	1.02
T _g (°C)	80

Synthesis Procedure:

ω -hydroxy terminated polystyrene was prepared by living anionic polymerization using OH protected initiator.

Characterization:

The molecular weight and polydispersity index of this polymer were determined by size exclusion chromatography (SEC) using a Varian liquid chromatograph equipped with a UV and refractive index detector. Polymer functionality was determined by titration with NaOH solution using phenolphthalein as the indicator.

Thermal analysis:


Thermal analysis of the samples was carried out using a differential scanning calorimeter (TA Q100) at a heating rate of 10°C/min. The inflection glass transition temperature (T_g) has been considered.

Solubility:


Polymer is soluble in toluene, THF, CHCl₃ and can be precipitated in water and cold methanol.

SEC of Sample:
Sample ID: P18845-SOH

Concentration (mg/mL)	15.8481
Sample dn/dc (mL/g)	0.1850
Method File	PS80K-aug.st 12/2014/0000.vcm
Column Set	3x PL 11136300
Solvent	THF



Sample	Mn	Mw	Mp	Mw/Mn	IV
P18845-SOH_01.vdt	8,512	8,744	8,755	1.027	0.0722



DSC thermogram for the sample:

