


**Sample: Poly(Styrene–co–Methyl Methacrylate),
 α -Hydroxy, ω -TEMPO-moiety terminated random copolymer**

Sample # P20252E-SMMAranOHT

Structure:




Composition:

$M_n \times 10^3$ (g/mol)	M_w/M_n (PDI)
17.7	1.26
Polystyrene content: 60 mol %	

Synthesis:

Hydroxy-terminated poly(styrene–co–methyl methacrylate) was prepared by stable free radical polymerization at 135 °C. The reaction scheme is shown below:



Characterization:


The molecular weight and polydispersity index (PDI) of the product was determined by size exclusion chromatography (SEC), using polystyrene as a standard. The ratio between polystyrene and poly(methyl methacrylate) in PS-PMMA copolymer was calculated from ^1H NMR spectroscopy by comparing the peak area of the PS phenyl protons at 6.5–7.3 ppm and the peak area of PMMA methyl protons at 2.6–3.6 ppm.

Solubility:

Poly(styrene–co–methyl methacrylate) is soluble in THF, DMF, toluene, and chloroform. It precipitates from methanol and hexanes.

^1H NMR spectrum of the copolymer in CDCl_3 :


1H NMR (500 MHz, CDCl_3): P20252-E (S:MMA=60:40)



SEC elugram of the copolymer:

Sample IDP20252E-SMMAranOHT

Concentration (mg/mL)	1.6159
Sample dv/dc (mL/g)	0.1300
Method File	P380K-June30-2015-0000.vsm
Column Set	3x PL 1113-6300
Solvent	THF



Sample	MW Number Average (Da)	MW Weight Average (Da)	MW at Peak (Da)	Polydispersity	Intrinsic Viscosity (dL/g)
P20252E_01.vdt	17,704	22,302	18,170	1.260	0.7138